Magnetic Detachment and Plume Control in Escaping Magnetized Plasmas

IEPC-2009-262

Presented at the 31st International Electric Propulsion Conference, University of Michigan • Ann Arbor, Michigan • USA September 20 – 24, 2009

> Paul F. Schmit¹ and Nathaniel J. Fisch² Princeton University, Princeton, NJ, 08540, USA

Abstract: The model of two-fluid, axisymmetric, ambipolar magnetized plasma detachment from thruster guide fields is extended to include plasmas with non-zero injection angular velocity profiles. Certain plasma injection angular velocity profiles are shown to narrow the plasma plume, thereby increasing exhaust efficiency. As an example, we consider a magnetic guide field arising from a simple current ring and demonstrate plasma injection schemes that more than double the fraction of useful exhaust aperture area, more than halve the exhaust plume angle, and enhance magnetized plasma detachment.

I. Introduction

THIS paper was published in the Journal of Plasma Physics, vol. 75, part 3, pp. 359-371 in 2009. Hard copies will be available during the David Fearn Memorial Poster Reception, which will be held in the Power Center Lobby from 6:00 - 7:30 pm on Monday, September 21, 2009.

¹ Ph.D. candidate, Department of Astrophysical Sciences, pschmit@princeton.edu. ²Professor, Department of Astrophysical Sciences, fisch@princeton.edu.